

LD Biopharma, Inc. 7384 Trade Street, Suite B San Diego, CA 92121 Tel: 858-876-8266 http://www.ldbiopharma.com

- PRODUCT DATA SHEET -

Name of Product: Recombinant YFP-HBxAg Fusion Protein

Catalog Number: VRP-3455

Manufacturer: LD Biopharma, Inc. USA

Introduction

HBxAg gene encodes a 17-kDa viral protein that plays an essential role in the HBV replication cycle. It was recently determined that a key function of HBxAg is to promote the degradation of the cellular structural maintenance of chromosomes 5/6 complex (Smc5/6). Smc5/6 directly binds DNA, and has been shown to topologically entrap DNA plasmids. As such, HBxAg is a potential therapeutic target since it promotes the degradation of the hepatocyte Smc5/6 complex that inhibits HBV transcription.

Full-length HBxAg cDNA (153aa, derived from Adw2 subtype) was constructed with codon optimization gene synthesis and expressed with YFP protein as N-terminal (YFP; 256aa) fusion partner in *E.coli* as inclusion bodies. The final product was refolded using our unique "temperature shift inclusion body refolding" technology and chromatographically purified.

Gene Symbol: HBxAg

Accession Number: AAK97176.1

Species: Viral

Size: 50 µg / Vial

Composition: 1.0 mg/ml, sterile-filtered, in 20 mM pH 8.0 Tris-HCl Buffer, with

proprietary formulation of NaCl, KCl, EDTA, Sucrose, DTT and

others.

Storage: In Liquid. Keep at -80°C for long term storage. Product is stable

at 4 °C for at least 30 days.

Key References

Mark Feitelson, et al. Hepatitis B virus x antigen in the pathogenesis of chronic infections and the development of HCC. American Journal Of Pathology 150(4):1141-57. (1997)



LD Biopharma, Inc.

7384 Trade Street, Suite B San Diego, CA 92121 Tel: 858-876-8266 http://www.ldbiopharma.com

Kornyeyev D, et al., Spatiotemporal analysis of hepatitis B virus X protein in primary human hepatocytes. J Virol 93:e00248-19. https://doi.org/10.1128/JVI.00248-19.(2019)

Decorsiere A, et al., Hepatitis B virus X protein identifies the Smc5/6 complex as a host restriction factor. Nature 531:386 -389. https://doi.org/10.1038/nature17170.(2016)

Applications

- 1. May be used for in vitro HBxAg protein mediated signal regulation for hepatocytes study using intracellular delivery of recombinant YFP-HBxAg protein with protein delivery reagent such as ProFectin.
- 2. May be used for HBxAg protein-protein interaction assay.
- 3. May be used as specific substrate protein for HBxAg specific kinase, and ubiquitin (Sumo pathway) related enzyme functional screening assays.
- 4. Potential therapeutic protein, by blocking HBxAg / host target protein interactions may be a useful strategy to control HBV replication or preventing liver cancer.
- 5. As native HBxAg antigen for its specific antibody production.

Quality Control

Purity: > 92 % by SDS-PAGE.

YFP protein: $\mathbf{Ex \lambda} = 517$ nm, and $\mathbf{Em \lambda} = 530$ nm.

Recombinant YFP- HBxAg Fusion Protein Sequence (45.8 kD)

MKHHHHHHQVSKGEELFTGVVPILVELDGDVNGHKFSVSGEGEGDATYGKLTLKLLCTTGKLPV PWPTLVTTLGYGVQCFARYPDHMKQHDFFKSAMPEGYVQERTIFFKDDGNYKTRAEVKFEGDTL VNRIELKGIDFKEDGNILGHKLEYNYNSHNVYITADKQKNGIKANFKIRHNIEDGGVQLADHYQ QNTPIGDGPVLLPDNHYLSYQSALFKDPNEKRDHMVLLEFLTAAGITEGMNELYK**GSENLYFQG**

EFAARLYCQLDPSRDVLCLRPVGAESRGRPFSGPLGTLSSPSPSAVPADHGAHLSLRRLPVCAF SSAGPCTLRFTSARCMETTVNAHQILPKVLHKRTLGLSAMSTTDLEAYFKDCVFKDWEELGEEI RLKVFVLGGCRHKLVCAPAPCNFFTSA