

LD Biopharma, Inc. 9924 Mesa Rim Road Suite B San Diego, CA 92121 Tel: 858-876-8266 http://www.ldbiopharma.com

- PRODUCT DATA SHEET -

Name of Product:Recombinant Human Cripto-1 ProteinCatalog Number:hRP-0544Manufacturer:LD Biopharma, Inc.

Introduction

Human teratocarcinoma-derived growth factor 1 (Cripto-1) gene encodes an epidermal growth factor-related protein that contains a cripto, FRL-1, and cryptic domain. The encoded protein is an extracellular, membrane-bound signaling protein that plays an essential role in embryonic development and tumor growth. Mutations in this gene are associated with forebrain defects. Recent data indicated that human Cripto-1 protein plays a role in the determination of the epiblastic cells that subsequently give rise to the mesoderm.

Full-length mature human Cripto-1 cDNA (31-150aa, isoform-1, derived from BC067844) was constructed with codon optimization with a small T7-His-TEV cleavage site Tag (29aa) fusion at its N-terminal. This protein was expressed in E. coli as inclusion bodies, refolded using our unique "temperature shift inclusion body refolding" technology and chromatographically purified.

Gene Symbol:	Cripto-1 (TDGF1; CR; CRGF)
Accession Number:	NP_003203
Species:	Human
Size:	20 µg / Vial
Composition:	0.2 mg/ml, sterile-filtered, in 20 mM pH 8.0 Tris-HCl Buffer, with proprietary formulation of NaCl, KCl, EDTA, arginine, DTT and Glycerol.
Storage:	In Liquid. Keep at -20° C for long term storage. Product is stable at 4 °C for at least 30 days.

Key References

De Luca, A., et al., *Expression and functional role of CRIPTO-1 in cutaneous melanoma*.



LD Biopharma, Inc. 9924 Mesa Rim Road Suite B San Diego, CA 92121 Tel: 858-876-8266 http://www.ldbiopharma.com

Br. J. Cancer 105 (7), 1030-1038 (2011)

Brandt,R., et al., *Identification and biological characterization of an epidermal growth factor-related protein: cripto-1*. J. Biol. Chem. 269 (25), 17320-17328 (1994)

Watanabe,K. et al., *Intercellular transfer regulation of the paracrine activity of GPIanchored Cripto-1 as a Nodal co-receptor*. Biochem. Biophys. Res. Commun. 403 (1), 108-113 (2010)

Applications

- 1. May be used for in vitro Cripto-1/Nodal signaling pathway regulations study for mesoderm differentiation.
- 2. Potential biomarker protein for various cancer diagnostic developments.
- 3. May be used as antigen for specific antibody production.

Quality Control

- 1. Purity: > 90% by SDS-PAGE.
- 2. Activity: Measured by its binding ability in a functional ELSA. Immobilized recombinant human Nodal protein at 2ug/ml (100ul/well) can bind rhCripto-1 with a linear range of 0.5-100 ng/ml.

Recombinant Protein Sequence

MASMTGGQQMGRGHHHHHHGNLYFQGGEFLGHQEFARPSRGYLAFRDDSIWPQEEPAIRPRSSQ RVPPMGIQHSKELNRTCCLNGGTCMLGSFCACPPSFYGRNCEHDVRKENCGSVPHDTWLPKKCS LCKCWHGQLRCFPQAFLPGCD