

LD Biopharma, Inc. 9924 Mesa Rim Road Suite B San Diego, CA 92121 Tel: 858-876-8266 http://www.ldbiopharma.com

### - PRODUCT DATA SHEET -

Name of Product: Recombinant Human PRPS2 Protein

**Catalog Number:** hRP-1415

**Manufacturer:** LD Biopharma, Inc.

### Introduction

Human PRPS2 gene encodes a phosphoribosyl pyrophosphate synthetase that plays a central role in the synthesis of purines and pyrimidines. The encoded protein catalyzes the synthesis of 5-phosphoribosyl 1-pyrophosphate from ATP and D-ribose 5-phosphate. Recent data indicated that PRPS2 expression was tightly controlled by Myc protein activities and plays a critical role in oncogenesis.

Full-length human PRPS2 (317aa, Isoform-2, derived from BC110875) gene was constructed with 29 aa N-terminal T7 / His / TEV cleavage site Tag and expressed in E.coli as inclusion bodies.. The final product was refolded using our unique "temperature shift inclusion body refolding" technology and chromatographically purified.

**Gene Symbol:** PRPS2 (PRSII)

Accession Number: NP\_002756

**Species:** Human

Size:  $50 \mu g / Vial$ 

**Composition:** 1.0 mg/ml, sterile-filtered, in 20 mM pH 8.0 Tris-HCl Buffer, with

proprietary formulation of NaCl, KCl, Sucrose and DTT.

**Storage:** In Liquid. Keep at -80°C for long term storage. Product is stable

at 4 °C for at least 30 days.

## **Key References**

John T. Cunningham, et al., *Protein and Nucleotide biosynthesis are coupled by a single rate-limiting enzyme, PRPS2, to drive cancer.* Cell 157, 1088 – 1103, May 22. (2014)

Mannava S, et al., Direct role of nucleotide metabolism in C-MYC-dependent proliferation of melanoma cells. Cell Cycle 7 (15), 2392-2400 (2008)



LD Biopharma, Inc. 9924 Mesa Rim Road Suite B San Diego, CA 92121 Tel: 858-876-8266 http://www.ldbiopharma.com

# **Applications**

- 1. May be used for in vitro PRPS2 mediated *myc signaling* regulation study for cancer cell metabolic rate controlling by intracellular delivery of this protein with "ProFectin" reagents.
- 2. May be used for mapping protein-protein interaction.
- 3. May be used as specific substrate protein for kinase and ubiquitin (Sumo pathway) related enzyme functional screening assays.
- 4. As antigen for specific antibody production.

# **Quality Control**

Purity: > 90% by SDS-PAGE.

## **Recombinant Protein Sequence**

MASMTGGQQMGRGHHHHHHGNLYFQGGEFPNIVLFSGSSHQDLSQRVADRLGLELGKVVTKKFS
NQETSVEIGESVRGEDVYIIQSGCGEINDNLMELLIMINACKIASSSRVTAVIPCFPYARQDKK
DKSRAPISAKLVANMLSVAGADHIITMDLHASQIQGFFDIPVDNLYAEPAVLQWIRENIAEWKN
CIIVSPDAGGAKRVTSIADRLNVEFALIHKERKKANEVDRMVLVGDVKDRVAILVDDMADTCGT
ICHAADKLLSAGATKVYAILTHGIFSGPAISRINNAAFEAVVVTNTIPQEDKMKHCTKIQVIDI
SMILAEAIRRTHNGESVSYLFSHVPL